

Association of *Calpain* Gene Polymorphism and Type Traits in Iranian One-hump Camels

Elham Rezvannejad^{1*}, Seyed Azim Mousavizadeh², Eisa Jorjani³, Mohammad Poortorogh⁴

1. Department of Biotechnology, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran

2. Department of Animal Science, Faculty of Agriculture, Ferdowsi University of Mashhad, Mashhad, Iran

3. Department of Biology, Faculty of Science, GonbadKavous University, GonbadKavous, Iran

4. Institute of Technical and Vocation of Agriculture Jahad, Hormozgan, Iran

Abstract

The present study investigated the calpain gene diversity in Iranian one-hump camels (120 camels) using PCR-RFLP method. Three genotypes AA, AG and GG were observed with the frequencies of 0.02, 0.19 and 0.78, respectively. The frequencies of A and G alleles were 0.12 and 0.88, respectively. The Chi-squared test didn't show significant deviations from Hardy-Weinberg equilibrium ($P < 0.05$). Association between the genotypes and type traits including body length, heart girth, height at withers, height at back and leg girth was modeled using the GLM procedure of the SAS program. There was a significant association between calpain genotypes and biometric traits, except for the body length ($P < 0.05$). The GG genotype had a greater height at back (ones of type traits) compared to other genotypes. The type traits including height at withers, heart girth and abdomen girth for GG genotype were significantly higher than the AA genotype but had no significant difference with the GA genotype ($P < 0.05$). The results suggested that calpain gene polymorphism may be a potential genetic marker for improving the growth-related traits in camels. More studies are required to confirm the effects of calpain gene on growth traits in camels.

Key words: Biometric trait; Calpain gene; Camel; PCR-RFLP; Polymorphism

Introduction

Since ancient times, domestic even-toed ungulates have played pivotal roles for man, being exploited for meat and milk, for fiber production, as beasts of burden for transport in agricultural/ rural oriented community; they were even worshipped. These animals have served for traditional

technologies since the very early era of domestication (1).

Camel belongs to the order Artiodactyla, sub-order Tylopod, and family of Camelidae. The genus *Camelus* consists of *Camelus dromedarius* (one hump) and *Camelus bactrianus* (two humps). Both species have been domesticated and reared for producing milk and meat. The meat of camels has been used as food in various parts of world for many centuries (2).

Genetic improvement has been considered as an important factor in the devel-

* Elham Rezvannejad, PhD

Department of Biotechnology, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran

Email: Rezvannejad2002@yahoo.com

Submission Date: 10 Jul. 2016 • Acceptance Date: 25 Aug. 2016

opment of domestic animal productions. Identifying the associated genes with economic traits as well as understanding their interactions with the environment or the other genes might be useful for applications of marker-assisted selection in the commercial animal population (3). Association of several polymorphic regions in candidate genes with economic traits has been considerably investigated in different animal species as cattle (4-7), goat (8,9), sheep (10,11), poultry (12,13) and camel (14).

The calpain-calpastatin system (CCS) includes a family of calcium-dependent neutral proteases. It is more likely to be found in most of the animal tissues (15). The first protein of CCS (m-calpain) was found in 1976 and μ calpain and calpastatin were recognized several years later. The activity of calpain depends on Ca^{2+} , as this enzyme plays a significant role in the growth of muscle and slaughtered meat tenderness through proteolysis of myofibrils (16). So far, the sequencing of regulatory subunit of calpain gene has been done by many researchers, all emphasizing the polymorphism of this region. In addition, other studies investigated the effects of this polymorphism on growth traits.

Chung et al. (17) amplified some parts of calpain III gene in sheep genome and identified the point mutations in this region using PCR-SSCP method. Furthermore, Chung et al. (18) has found that there is a meaningful difference between fat around the hips, kidneys and heart of the different genotypes for calpain gene. They also studied this region in Angus breed cattle and its relationship with slaughtered meat

tenderness and carcass traits. In addition, the effects of this gene on beef quality traits have been discussed by Esmailizadeh et al. (19). Human studies have associated calpain-10 variants with type 2 diabetes and metabolic traits (20).

The calpain alleles are easily detected by PCR amplification (17). Therefore, calpain gene was investigated as a candidate gene to identify quantitative trait loci (QTL) affecting slaughtered meat tenderness and body weight traits in many species of animals.

Determination of gene polymorphism is important in breeding of farm animals (21,22). GhasemiMeymandi et al. (23) investigated genetic variation within Iranian indigenous *Camelus dromedarius* in northern Kerman province using five microsatellite markers. The results of the study indicated that the *Camelus dromedarius* in the North of Kerman province populations had a relatively high genetic variation resulting in adaptation with drastic environmental changes of Kerman province. In another study, Ghasemi Meymandi et al. (24) investigated individual's assignment of the Camel Populations of North of Kerman Province using 7 different methods. Among the methods based on the likelihood, the Rannala and Mountain method and among the methods based on the genetic distance, Nei DA method showed the highest accuracy. Furthermore, Ghasemi Meymandi et al. (25) analyzed genetic structure of *Camelus dromedarius* using PCA and Hierarchical clustering methods and concluded that all samples could be mainly regrouped into three main clusters and most suitable number of genetic

groups in dataset was explained and a similarity of geographical distribution was in good accordance with founded genetic relationships in this study.

However, no study concerning the polymorphism of calpain gene in Iranian camel breed (one hump) and its association with body measurements has been conducted yet. Thus, this study aimed to genotype the calpain gene in Iranian camel breed (one hump) for investigating the possibility of its correlation with body measurements.

Materials and Methods

Animals and phenotypic traits

One hundred and twenty camels were randomly selected from different herds in Hormozgan province, Iran. Blood samples (approximately 8 ml) were collected from jugular veins of the camels in EDTA-coated tubes and stored at -20°C until DNA extraction.

In Table 1, the measured type traits including length of body, heart girth, abdomen girth, height at withers and height at back are shown. Also, sex, age, type of birth and pregnancy state of animals were recorded.

Table 1: Overall mean, number of observations and phenotypic standard deviation for measured traits

trait	Number	Mean (cm)	Standard deviation
length of body	120	148.25	0.10
heart girth	120	172.43	0.13
abdomen girth	120	165.56	0.15
height at back	120	179.21	0.09
height at withers	120	219.75	0.13

Genomic DNA extraction

Total genomic DNA was extracted from blood samples using Accuprep Genomic DNA Extraction Kit, Cat No: K-3032

according to the manufacturer protocol. Quantity of the extracted DNA was measured according to spectrometry method using Nano-drop and 2000 spectrophotometer of USA Thermo Company and its quality was checked on 1% Agarose gel.

PCR Primers and Amplification

Primers for PCR amplification of calpain were designed with regard to the reported bovine CAPN1 sequences (Gen Bank AF252504), because of unavailability of sequence of this gene in camels and also Camelidae family. 120 camels were genotyped using the PCR-RFLP method. A 787 bp fragment was amplified with the primers of 5'-AGCGCAGGGACCCAGTGA-3' (forward) and 5'-TCCCCTGC-CAGTTGTCTGAAG-3' (reverse) using the following thermal cycle: an initial hot start at 95°C for 5 min followed by 35 cycles, denaturation at 95°C for 45s, annealing at 62°C for 45s, extension at 72°C for 50s and a final extension for 7 min at 72°C after 35 cycles. PCR reactions were performed in a total volume of 25µl containing 2.5µl of standard PCR buffer, 1.5mM MgCl₂, and 0.5µM of each primer, 2µM dNTP, 0.2 U Taq DNA polymerase and 120ng of template DNA. The PCR products were analyzed by electrophoresis on a 1% agarose gel and the gel was stained with ethidium bromide. In addition, the marker size was used for confirming the accuracy of desired fragment. 16 ng of PCR products were digested using the restriction enzyme *Ava*II (2U). Restriction site of this enzyme is 5'---G/GWCC---3' and 3'---CCWG/G---5'. The digested fragments were analyzed by electrophoresis in

2% agarose gel, stained with ethidium bromide and photographed under UV light.

Statistical analyses

The observed allele frequency and the Hardy-Weinberg equilibrium were analyzed using the PopGene32 software. For the association study, the traits were analyzed using general linear model (GLM) of Statistical Analysis System (SAS Institute, 2004) program least squares means of genotypes were compared by the Duncan test. The applied linear model was as follow:

$$Y_{ijklns} = \mu + \text{Sex}_i + \text{Age}_j + \text{Type}_k + \text{Pregnancy}_l + \text{genotype}_n + e_{ijklns}$$

Where Y_{ijklns} is the trait of interest, μ is the general mean, Sex_i is the fixed effect of i th sex ($i=1,2$), Age_j is the fixed effect of j th age ($j=1, \dots, 4$), Type_k is the fixed effect of k th type of birth ($k=1,2$), Pregnancy_l is the fixed effect of l th Pregnancy state ($l=1,2$), Genotype_n is the fixed effect of n th genotype ($n=1,2,3$), and e_{ijklns} is the random error.

Results

Genotype and allele frequencies

A 787bp amplicon of exon 14 comparable to that of the bovine amplicon was amplified by PCR for camel calpain gene. The PCR products were electrophoresed on 1 % agarose gel shown in Figure 1. The PCR-RFLP analysis displayed three geno-




Figure 1: Electrophoresis of PCR products for 14 exon region of calpain gene

types including AA, AG and GG.

The observed number, observed genotype frequencies, the expected frequency, the expected number, Chi-square (χ^2) and observed allele frequencies in the exon 14 of calpain gene in Iranian camel breed (one hump) have been shown in Table 2. The allelic frequencies were 0.878 and 0.122 for G and A, respectively. Genotype GG showed maximum observed frequency (78.33%) in the herd. Heterozygous genotype AG showed lower frequency (19.17%) than the genotype GG while the genotype AA had lowest frequency (2.5%) in the testing herd. To evaluate Hardy-Weinberg equilibrium, Chi-squared test was applied and the results have been shown in Table 2. Differences of observed genotypes and expected genotypes were not significant according to the calculation of $\chi^2 = 1.247$, suggesting that the studied population for calpain gene is in the Hardy - Weinberg equilibrium.

Table 2: Observed number, observed genotype frequencies, expected frequency, expected number, Chi-square (χ^2) and observed allele frequencies of exon 14 of calpain gene in camel

genotype	Observed genotype number	observed genotype frequencies	expected genotype frequency	expected genotype number	Chi-square χ^2 (O-E) ² /E	observed allele frequencies	
						G	A
AA	3	2.5	1.425	1.71	0.97		
AG	23	19.17	21.3	25.57	0.26	0.878	0.122
GG	94	78.33	77.275	92.72	0.0179		
total	120	100	100	120			

Having observed and expected heterozygosity, Nei expected heterozygosity and the average heterozygosity of calpain gene in the studied herd are given in Table 3. Heterozygous level is considered as one of the indicators of introducing the genetic variations in a population. Average heterozygosity, Nei heterozygous and observed heterozygosity were 0.193, 0.214 and 0.214, respectively.

Table 3: Observed and expected heterozygosity, Nei expected heterozygosity and the average heterozygosity of exon 14 of calpain gene in camels

gene	Observed heterozygosity	expected heterozygosity	Nei expected heterozygosity	average heterozygosity
calpain	0.193	0.215	0.214	0.214

Association between genotypes and type traits

There were significant detectable effects of calpain genotypes on the measured type traits except length of body in camels ($P < 0.05$) (Table 4). The height at withers, heart girth and abdomen girth traits for GG genotype were higher than those for the AA genotype, but there was no significant difference in these traits between GG and AG genotypes ($P > 0.05$). Also, the height at back for GG genotype was significantly higher than that of the AG and AA genotypes ($P < 0.05$).

Table 4: Least square means and standard errors of the length of body, heart girth, abdomen girth, height at withers and height at back determined for genotypes of calpain polymorphism

Trait (cm)	Genotypes		
	AA	AG	GG
length of body	147.33±0.12 ^a	148.46±0.11 ^a	150.74±0.14 ^a
height at withers	164.33±0.17 ^b	174.12±0.13 ^a	174.29±0.11 ^a
height at back	157.00±0.11 ^c	166.33±0.12 ^b	184.97±0.09 ^a
heart girth	155.00±0.09 ^b	182.54±0.09 ^a	180.32±0.08 ^a
abdomen girth	174.33±0.15 ^b	225.29±0.11 ^a	218.92±0.13 ^a

Discussion

Calpain has been considered as a candidate gene for carcass performance and meat quality traits in farm animals. In livestock, variations in the calpain gene have been detected in the cattle, sheep and pigs, but not in camels. Page et al. (26) recognized marker calpain1 (an adenosine/guanosine (A/G) substitution) in the exon 14 of bovine calpain1 gene. This marker has been suggested to be associated with meat tenderness (26-30).

In the present study, three different genotypes (AA, AG and GG) were found in calpain1 locus. Similar result for calpastatin locus was observed in Sistani cattle by FakhrKazemi et al. (31). FakhrKazemi et al. (31) carried out a study on polymorphism of calpastatin gene for Sistani cattle and MM, MN, NN genotypes with frequencies of 51.86, 32.27 and 4.87 with respect to the calpastatin gene of cattle were reported. M and N allele frequencies were given as 76.4 and 23.6%, respectively. Low heterozygosity of calpastatin gene of cattle was obtained from a closed herd.

A high degree of calpain polymorphism has also been reported by Azari et al. (32), Bahrapour et al. (15), Mohammadabadi (33) and Arora et al. (34) in different breeds of sheep. Azari et al. (32) observed 3 different genotypic patterns (G1, G2 and G3)

with the frequencies of 0.082, 0.892 and 0.027 in Iranian Dalagh sheep. The results of their research conformed to earlier reports of Tahmoorespour (35) who reported three genotypes (AA, AB and BB) with allele frequencies of 0.56 and 0.44 for A and B alleles respectively in Baluchi sheep. Bahrampour et al. (15) also reported 3 genotypes MM, MN and NN with frequencies of 0.69, 0.20 and 0.11 respectively in Kermani sheep. Mohammadabadi (33) demonstrated that 3 genotypes MM, MN and NN had frequencies of 0.67, 0.25 and 0.08, respectively in Sanjabi sheep. Arora et al. (34) studied 11 Indian sheep breeds and studied the allele frequencies of 0.603 and 0.397 for A and B alleles respectively and frequencies of 0.388, 0.429 and 0.183 for AA, AB and BB genotypes, respectively. Furthermore, the observed and expected heterozygosity were estimated as 43.5 and 56.9%, respectively in this research.

These results were consistent with those obtained in the present study in camel. Accordingly, there is no selective advantage or disadvantage of a particular genotype in this population and Hardy-Weinberg equilibrium is established in this camel herd. Moreover, moderate and high heterozygous levels of calpain gene demonstrated good genetic variations based on obtained results of average heterozygosity, Nei heterozygous and observed heterozygosity in the studied population. This may be explained by the conservation and breeding strategies, which have been carried out.

Many studies have reported a relationship between polymorphism of calpain gene and quality of carcass traits while some researchers have investigated the re-

lationships between polymorphism of this gene and growth traits in cattle and sheep (36-38). Development of skeletal muscle depends on three main factors, these factors involve protein synthesis of muscles and decomposition of muscle proteins and muscle cells. Increased skeletal muscle growth could be associated with decreased protein decomposition of muscle caused by calpain system activity. Calpain gene is introduced as an effective candidate gene in meat production (15,33).

The results of our research on the association between calpain gene and type traits in camel were verified with previous reports in other species on the body weight traits. Cheong et al. (3) examined the association between polymorphism of the micro-calpain gene and carcass traits in Korean cattle. The results revealed that there was a significant association between the 3'UTR and marbling score ($P < 0.02$). Furthermore, Chung et al. (39) investigated the association between ovine calpain gene polymorphism and body weight traits. Body weight was recorded at birth, weaning and post-weaning. Calpain 3 genotypes of CAPN31112 segment were associated with birth weight ($P < 0.01$) and a dominant gene effect was observed. Also, associations between m-calpain/HhaI polymorphism and growth traits in Chinese indigenous cattle breeds were studied by Zhang and Li (38). The m-calpain/HhaI polymorphism was associated with body weight, withers height, and body length for 6 months ($P < 0.05$ or $P < 0.01$), body length for 18 months ($P < 0.05$), and body length and heart girth for 24 months ($P < 0.01$) in Nanyang cattle.

The public combining ability of a gene is caused by additive effects in general, but for a specific combining ability of a gene the dominant effects and epistasis are important (40). In this study a dominant effect was recognized for height at withers, heart girth and abdomen girth traits. The effect may shape the specific combining ability. No dominant effects were found for height at back. The results suggest that type traits may be related to genetic factors as well as environmental factors. Some environmental factors were confirmed to be significant in this study.

Linear type classification is an important tool in decision making as it focuses on the selection of animals that should have a longer herd life while expressing their productive and reproductive potentials based on their morphologic traits (41). Previous researches have indicated the usefulness of linear type traits as predictors of body weight (42,43), health (44) and fertility (44,45) in dairy cattle. The correlations between type traits and performance were estimated as moderate to high (46,47).

Janssens and Vandepitte (48) estimated genetic parameters for seven body measurements and 13 linearly scored traits in

three sheep breeds. Heritability of measured traits was in the range of 0.26–0.57 and genetic correlations were high between these traits. Also, Ishag et al. (14) estimated weights of camels using the Boue formula (1949) with regard to type traits. Moreover, knowledge of association between polymorphisms and performance can increase the accuracy of selection.

Results of this research showed that calpain marker had a detectable effect on type traits in camels. However, with regard to high correlation between type traits and body weight traits, calpain gene is considered as a main economic gene due to its relation with productive traits in camels and it may explain variations of either body weight or growth. The polymorphism may also be useful in marker-assisted selection for body weight.

Acknowledgements

This research was carried out with the financial support project (No. 7/95/2359/ص) by Graduate University of Advanced Technology, Kerman.

References

- Barazandeh A, Mohammadabadi MR, Ghaderi M, Nezamabadipour H. Genome-wide analysis of CpG islands in some livestock genomes and their relationship with genomic features. *Czech Journal of Animal Science* 2016;61:487-495.
- Tahmoospour M, Shojaei M. Partial sequencing and investigation of leptin and calpain, candidate genes for meat quality in camel. *Journal of Camel Practice and Research* 2013;20(2):277-282.
- Cheong H, Yoon D, Park BL, et al. A single nucleotide polymorphism in CAPN1 associated with marbling score in Korean cattle. *BMC Genetics* 2008;9:33-40.
- Kharrati Koopaei H, Mohammad Abadi MR, Ansari Mahyari S, et al. Effect of DGAT1 variants on milk composition traits in Iranian Holstein cattle population. *Animal Science Papers & Reports* 2012;30(3):231-239.
- Ebrahimi Z, Mohammadabadi MR, Esmailzadeh A, Khezri A. Association of PIT1 gene and milk protein percentage in Holstein cattle. *Journal of Livestock Science and Technologies* 2015a;3(1):41-49.
- Ebrahimi Hoseinzadeh Z, Mohammadabadi MR, Esmailzadeh Koshkueh A, et al. Association of PIT1 gene with milk fat percentage in Holstein cattle. *Iranian Journal of Applied Animal Science (IJAS)* 2015b;5(3):575-582.
- Pasandideh M, Mohammadabadi MR, Esmailzadeh AK, Tarang A. 2015. Association of bovine PPARG1A and OPN genes with milk production and composition in Holstein cattle. *Czech Journal of Animal Science* 2015;60(3):97-104.
- Shamsalddini S, Mohammadabadi MR, Esmailzadeh AK. Polymorphism of the prolactin gene and its effect on fiber traits in goat. *Russian Journal of Genetics* 2016;52(4):1-5.
- Moghadaszadeh M, Mohammadabadi MR, Esmailzadeh Koshkueh A. Association of Exon 2 of BMP15 Gene with the Litter Size in the Raini Cashmere Goat. *Genetics in the 3rd Millennium* 2015;13(3):4062-4067.
- Shojaei M, Mohammad Abadi MR, Asadi Fozzi M, et al. Association of growth trait and Leptin gene polymorphism in Kermani sheep. *Journal of Cell and Molecular Research* 2011; 2(2): 67-73.
- Zamani P, Akhondi M, Mohammadabadi MR. Associations of Inter-Simple Sequence Repeat loci with predicted breeding values of body weight in Sheep. *Small Ruminant Research* 2015;132(3):123-127.
- Moazeni SM, Mohammadabadi MR, Sadeghi M, et al. Association of the melanocortin-3(MC3R) receptor gene with growth and reproductive traits in Mazandaran indigenous chicken. *Journal of Livestock Science and Technologies* 2016a;4(2):51-56.
- Moazeni, S.M., Mohammadabadi, M.R., Sadeghi, M., et al. Association between UCP Gene Polymorphisms and Growth, Breeding Value of Growth and Reproductive Traits in Mazandaran Indigenous Chicken. *Open Journal of Animal Sciences* 2016b;6:1-8.
- Ishag IA, Reissmann M, Peters KJ, Musa LMA, Ahmed MKA. Phenotypic and molecular characterization of six Sudanese camel breeds. *South African Journal of Animal Science* 2010;40(4):319-326.
- Bahrpour V, Mohammadabadi MR, Mirzaei, H.R., et al. Molecular analysis of calpastatin gene in Kermani sheep herds. *Journal of Agricultural Sciences and Natural Resources* 2008;15(4):124-131. (In Persian)
- Sensky PL, Parr T, Bardsley RG, Buttery PJ. Postmortem proteolysis in muscle and meat quality and phenotypic and genetic correlations for bovine post rigor calpastatin activity, intramuscular fat content, variable activity of the calpain proteolytic system. Meat and Livestock Commission, Loughborough, UK 2000.
- Chung HY, Davis ME, Hines HC. A DNA polymorphism of the bovine calpastatin gene detected by SSCP analysis. *Animal Genetics* 1999a;30(1):80-81.
- Chung HY, Davis ME, Hines HC, Wulf DM. Relationship of a PCR-SSCP at the bovine calpastatin locus with calpastatin activity and meat tenderness. *Ohio State University Research and Reviews: Beef and Sheep* 170. 1999b. <http://ohioline.osu.edu/sc170/sc170-3.html> [Verified 18 June 2004].
- Esmailzadeh AK, Morris CA, Cullen NG, et al. Genetic mapping of quantitative trait loci for meat quality and muscle metabolic traits in cattle. *Animal Genetics* 2011;42(6):592-9.
- Cheverud JM, Fawcett GL, Jarvis P, et al. Calpain-10 is a component of the obesity-related quantitative trait locus Adip1. *Journal of Lipid Research* 2010;51(5):907-913.
- Mohammadi A, Nassiry MR, Mosaferi J, Mohammadabadi MR, Sulimova GE. Distribution of BoLA-DRB3 allelic frequencies and identification of a new allele in the Iranian cattle breed Sistani (Bosindicus). *Russian journal of genetics* 2009;45(2):198-202.
- Mousavizadeh A, Mohammad Abadi, M, Torabi, A, et al. Genetic polymorphism at the growth hormone locus in Iranian Tali goats by polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP). *Iranian Journal of Biotechnology* 2009;7(1):51-53.
- GhasemiMeymandi M, Mohammadabadi MR, Esmailzadeh AK. Genetic variation of camels in the North of Kerman province using microsatellite markers. *Animal Production Research* 2015;4(1):35-45. (In Persian)
- Ghasemi Meymandi M, Mohammadabadi MR, Esmailzadeh A, Montazeri M. Assigning Individuals to the Camel Populations of North of Kerman Province Using Microsatellite Markers. *Modern Genetics Journal* 2016a;11(3):329-335. (In Persian)
- Ghasemi Meymandi M, Mohammadabadi MR, Montazeri M. Analyzing Genetic Structure of Camelus dromedarius Using PCA and Hierarchical clustering methods. *Journal of Agricultural Biotechnology* 2016c;8(3):83-96 (In Persian)
- Page BT, Casas E, Heaton MP, et al. Evaluation of single-nucleotide polymorphisms in CAPN1 for association with meat tenderness in cattle. *Journal of Animal Science* 2002;80:3077-3085.
- Page BT, Casas E, Quaas RL, et al. Association of markers in the bovine CAPN1 gene with meat tenderness in large crossbred populations that sample influential industry sires. *Journal of animal science* 2004;82(12):3474-3481
- Allais S, Levéziel H, Payet-Duprat N, et al. Effects of polymorphisms in the calpastatin and μ -Calpain genes on meat tenderness in three French beef breeds. *Journal of Animal Science* 2011;89(1):1-11.
- Casas E, White SN, Wheeler TL, et al. Effects of calpastatin and μ -calpain markers in beef cattle on tenderness traits. *Journal of Animal Science* 2006;84:520-525.
- Othman EO, Fawzia AZ, Ali AEG, Medhat RAE. Genetic polymorphism of three genes associated with milk trait in Egyptian buffalo. *Journal of Genetic Engineering and Biotechnology* 2011;9(1): 15-20.
- FakhrKazemi FM, Nasiri MR, FathiNajafi M, EftekhariShahroudi F, Khosravi F. Studying genetic polymorphism of calpastatin gene in Systani cows using RFLP method, Proceedings of the 2th National Congress of Animal Science and Aquaculture, Mashhad, Iran 2007.
- Azari MA, Dehnavi E, Yousefi S, Shahmohamdi L. Polymorphism of calpastatin, calpain and myostatin genes in native Dalagh sheep in Iran. *Slovak Journal of Animal Science* 2012;45(1):1-6.
- Mohammadabadi MR. Allelic diversity of calpastatin gene in Sanjabi sheep. *Journal of Cellular and Molecular Researches (Iranian Journal of Biology)* 2016;28(3):395-402. (In Persian)
- Arora RR, Yadav HS, Yadav DK. Identification of novel single nucleotide polymorphisms in candidate genes for mutton quality in Indian sheep. *Animal Molecular Breeding* 2014;4:1-5.
- Tahmoospour M. Study on the calpain and calpastatin gene polymorphisms by polymerase chain reaction-single strand conformation polymorphism and its relation to average daily gain in Iranian Baluchi sheep. Ph.D. Thesis, Ferdowsi University, Mashhad, Iran 2005.
- Corva PM, Soria L, Villarreal EL, et al. Association of polymorphisms on the CAPN1 and CAST genes with meat tenderness in beef cattle of Argentina. *Genetics and Molecular Biology* 2007;30(4):1064-1069.
- Soria LA, Corva PM, Sica A, et al. Effect of three single-nucleotide polymorphisms in CAPN1 gene on beef tenderness. *ArchivTierzucht* 2009;52(5):546-549.
- Zhang R, Li X. Association between IGF -IR, m-calpain and UCP-3 gene polymorphisms and growth traits in Nanyang cattle. *Molecular Biology Reports* 2011;38(3):2179-2184.
- Chung H, Choi B, Jang G, et al. Effect of variants in the ovine skeletal-muscle-specific calpain gene on body weight. *Journal of Applied Genetics* 2007;48(1): 61-68.
- Falconer DS, Mackay TFC. Introduction to quantitative genetics, 4th ed. Harlow, Essex, 1996;274-283.
- Kern EL, Cobuci JA. Factor Analysis of linear type traits and their relation with longevity in Brazilian Holstein cattle. *Asian Australia Journal of Animal Science* 2014;27(6):784-790.
- Koenen EPC, Groen AF. Genetic evaluation of body weight of lactating Holstein heifers using body measurements and conformation traits. *Journal of Dairy Science* 1998;81(6):1709-1713.
- Veerkamp RF, Brotherstone S. Genetic correlations between linear type traits, food intake, live weight and condition score in Holstein Friesian dairy cattle. *Animal Science* 1997;64(3):385-392.
- Pryce JE, Esslemont RJ, Thompson R, Veerkamp RF, Kossabati MA, Simm G. Estimation of genetic parameters using health, fertility and production data from a management recording system for dairy cattle. *Animal Science* 1998;66(3):577-584.
- Royal MD, Pryce JE, Woolliams JA, Flint APF. The genetic relationship between commencement of luteal activity and calving interval, body condition score, production, and linear type traits in Holstein-Friesian dairy cattle. *Journal of Dairy Science* 2002;85(11):3071-3080.
- Berry DP, Buckley F, Dillon P, Evans RD, Veerkamp RF. Genetic relationships among linear type traits, milk yield, body weight, fertility and somatic cell count in primiparous dairy cows. *Irish Journal of Agricultural and Food Research* 2004;43(2):161-176.
- Tapki I, Guzey YZ. Genetic and phenotypic correlations between linear type traits and milk production yields of Turkish Holstein dairy cows. *Greener Journal of Agricultural Sciences* 2013;3(11):755-761.
- Janssens S, Vandepitte W. Genetic parameters for body measurements and linear type traits in Belgian Bleu du Maine, Suffolk and Texel sheep. *Small Ruminant Research* 2004;54(1-2):13-24.